

Burn

By

Dr. Sajjad M Alhakeem

F.I.B.M.S (Urology)




Burn is an injury to the skin or other organic tissue primarily caused by heat or due to radiation, radioactivity, electricity, friction or contact with chemicals.

Types of burns

• Thermal. • Chemical. • Electrical • Radiation

Thermal (heat) burns occur when some or all the cells in the skin or other tissues are destroyed by:

- hot liquids (scalds)
- hot solids (contact burns)
- flames (flame burns).



Assessing depth

FIRST-DEGREE BURNS

- Redness, tenderness, and pain are the hallmarks of this injury.
- Blistering does not occur,
- Healing takes place after several days without scarring



Second-degree burns

- These burns appear pink, moist, and soft and are exquisitely tender when touched by a gloved hand.
- They heal in approximately 2-3 weeks, usually without scarring



Third-degree burns

- Third-degree burns are full-thickness burns that destroy both epidermis and dermis
- may be white, brown, or black and look 'leathery'. They do not blister and have no sensation
- skin grafting always is necessary to resurface the injured area



Fourth-degree burns


- cause full-thickness destruction of the skin and subcutaneous tissue,
- with involvement of the underlying fascia, muscle, bone, or other structures.
- These injuries require extensive debridement and complex reconstruction of specialized tissues and invariably result in prolonged disability



Assessing extent of burns

ADULTS RULE OF 9'S:

INFANTS RULE OF 5'S



- head = 20 %
- each arm = 10 %
- each leg = 20 %
- front of trunk = 10 %
- back of trunk = 10 %

Major burns: resuscitation

Prehospital first aid measures

- Ensure rescuer safety first.
- Remove the patient from the burning environment. If clothes are smoldering, apply cold water and remove them, unless adherent.
- Treat airway obstruction
- Provide high flow O₂
- Cover burns in clean sheets
- Continue O₂ and apply a hard cervical collar if there is any possibility of spinal injury



Analgesia

- Obtain IV access with two large peripheral cannula.
- Send blood: X-matching, FBC, , U&E, glucose, and coagulation.
- Provide analgesia (IV morphine titrated according to response).
- Provide an antiemetic .

Fluid resuscitation

- Give IV fluids. Start with isotonic crystalloid (eg 0.9 % saline) at 2–4mL of
- crystalloid per kg body weight per % body surface area burned, over the
- first 24hr following injury. Give half of this volume in the first 8hr.
- Check pulse, BP, and respiratory rate every 10–15min initially.
- Insert a urinary catheter and test the urine

PARKLAND FORMULA

$$4 \text{ mL} \times \% \text{ BSA} \times \text{weight (kg)} = \text{volume (mL)}$$

1/2 1/2
↓ ↓
first 8 hrs next 16 hrs

Burn Wound Management



- powder-free gloves because of the demonstrated toxicity of glove powders to tissue.
- Initial treatment of the burn wound involves cleansing the wound with saline
- Ruptured blisters are removed with scissors.
- After wound cleansing, cover with a topical antimicrobial dressing ,Topical antibiotics decrease microbial growth and reduce invasive infection.
- Systemic antibiotics may be indicated when cellulitis is evident in surrounding unburned tissue.

